



Com|Pl|i|t DX

ΠΛΗΡΟΦΟΡΙΕΣ

Εξεταζόμενος	-	Ημ. Συλλογής Δείγμ.	-
ΑΜΚΑ	-	Παραπέμπων	-
Ημερ. Γέννησης	-	Barcode	26xxxxxxGR
Τύπος Δείγμ. #1	ΙΣΤΟΣ ΕΓΚΛΕΙΣΜΕΝΟΣ ΣΕ ΜΠΛΟΚ ΠΑΡΑΦΙΝΗΣ	Ημερ. Παρ. Δείγμ.	-
Τύπος Δείγμ. #2		Ημερ. Αποτελ.	-
Κωδικός Δείγμ. #1	-	Τύπος όγκου	ΚΑΡΚΙΝΟΣ ΠΡΟΣΤΑΤΗ

Com.Pl.i.t. Dx (30 genes, 9 fusions) | Comprehensive Panel for Individualized Treatment

Αποτελέσματα και ερμηνεία*

Βιοδείκτης	Αποτέλεσμα	Θεραπείες με ισχυρή κλινική σημασία		Θεραπείες με πιθανή κλινική σημασία	Θεραπείες με πιθανή αντίσταση
		Εγκεκριμένες/ βάσει οδηγιών θεραπειές για την ένδειξη (Level A)	Θεραπείες σε ισχυρές μελέτες (Level B)	Εκτός ένδειξης θεραπείες (Level C)	
TP53	Exon 8 p.R273C VAF: 43%	-	-	-	-
Μικροδορυφορική Αστάθεια (MSI)	χωρίς μικροδορυφορική αστάθεια (MSS)	-	-	-	-

*Σημείωση: Το επίπεδο σημαντικότητας των παραλλαγών (Level of Evidence, LoE) (π.χ. A, B, C κλπ) βασίζονται στις οδηγίες για την αναφορά γενετικών παραλλαγών στον καρκίνο που δόθηκαν με κοινή συναίνεση των AMP, ACMG, ASCO και CAP. Για λεπτομερή περιγραφή των οδηγιών αυτών, ανατρέξτε στην Εικόνα 1.

^ Η ανάλυση συγκεκριμένων βιοδεικτών του παρόντος πολυγονιδιακού πάνελ πραγματοποιείται στο πλαίσιο διαπίστευσης κατά ΕΛΟΤ EN ISO 15189:2022 (βλ. ενότητα «Methodology»). Η διαπίστευση κατά ΕΛΟΤ EN ISO 15189:2022 αφορά αποκλειστικά τον αναλυτικό προσδιορισμό των εν λόγω βιοδεικτών. Οι ερμηνείες που σχετίζονται με την πρόβλεψη ανταπόκρισης σε θεραπεία δεν περιλαμβάνονται στο πεδίο διαπίστευσης.



Αρ. 822

Λεωφ. Σπάτων 52, 15344, Γέρακας, Αττική, αρ. Γ.Ε.ΜΗ.: 0007856001000 | info@genekor.com, www.genekor.com | Τηλ. (+30) 210 6032138 Fax. (+30) 210 6032148
Μια εταιρεία πιστοποιημένη με ISO 9001:2015 (Cert. No 041150049) και ISO/IEC 27001:2022 (Cert. No 048190009)

Γιώργος Νασιούλας, PhD Μοριακός Βιολόγος, Επιστημονικός Διευθυντής, ΑΜΚΑ:26025301255

Document Code: Δ21_Prostate_FFPE



Com|Pl|i|t DX

SAMPLE INFORMATION

Name	-	Date Sp. Extracted	-
ID/Medical ID	-	Req. Physician	-
Date of Birth	-	Report No	26xxxxxxGR
Material #1	PARAFFIN EMBEDDED TISSUE-BLOCK	Date Received	-
Material #2	-	Date of Report	-
Sample #1 ID	-	Tumor type	PROSTATE CANCER

Com.Pl.i.t. Dx (30 genes, 9 fusions) | Comprehensive Panel for Individualized Treatment

1. Clinically Significant Biomarkers*

Biomarker	Result	Therapies with strong clinical significance		Therapies with potential significance	Therapies with potential resistance
		Approved/ Standard of care therapies (Level A)	Therapies in well powered studies (Level B)	Off label/ Investigational therapies (Level C)	
TP53	Exon 8 p.R273C VAF: 43%	-	-	-	-
Microsatellite Instability (MSI)	Stable (MSS)	-	-	-	-

*Note: Variants' Level of Evidence (LoE) (e.g. A, B, C etc) are based on the Joint consensus recommendation of AMP, ACMG, ASCO and CAP for reporting genetic variants in cancer. For a detailed description of the recommendation please refer to Fig. 1

^ The analysis of selected biomarkers included in the present multigene panel is performed within the scope of accreditation according to ELOT EN ISO 15189:2022 (see «Methodology» section). Accreditation according to ELOT EN ISO 15189:2022 applies exclusively to the analytical determination of the selected biomarkers. Interpretations related to the prediction of therapeutic response are not included in the scope of accreditation.



Name

-

Report No

26xxxxxxGR

1. [Clinically Significant Biomarkers](#)
2. [Methodology](#)
3. [Quality Control Results](#)
4. [Important biomarkers findings](#)
5. [Immune Checkpoint inhibitors biomarkers](#)
6. [Interpretations for targeted therapies](#)
7. [Clinical Trials to consider](#)
8. [Appendix](#)
9. [Biomarker findings across all Levels of evidence \(A-D\)](#)
10. [References](#)



52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 6032138 Fax. (+30) 210 6032148
A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022(Cert. No 048190009)

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE



Name

-

Report No

26xxxxxxGR

2. Methodology

NGS analysis

DNA is extracted from the sample under investigation using the Qiaamp DNA FFPE Kit (Qiagen). A capture based targeted next generation sequencing (NGS) analysis was performed, using the Oncology 188-Gene Variant Assay (GenePlus) which is a qualitative test which enables the simultaneous detection of single nucleotide variants (SNVs), insertions and deletions (InDels), fusions and copy number variations (CNVs) as well as microsatellite instability (MSI) in a single workflow.

Sequencing was carried out on the DNBSEQ-T7 NGS platform (MGI). The analysis of 30 genes includes the entire exon regions and CNVs of 11 genes, partial exon regions of 19 genes and introns/promoters/fusion breakpoint regions of 9 genes. The test also reports on Microsatellite Instability (MSI) status.

Sequencing data are analyzed through bioinformatics pipeline for variant calling and interpretation using the Gene+Box data analysis and management system. Alteration- and tumor type-specific therapeutic implications are classified using the OncoKB™ Levels of Evidence system, which assigns clinical actionability to individual mutational events. For additional details about the OncoKB™ curation process, please refer to the version-controlled OncoKB™ Curation Standard Operating Procedure.

Genes Analyzed

30 Gene Alterations

AR	ATM*	ATR	BARD1	BRAF	BRCA1*	BRCA2*	BRIP1	CDK12	CHEK1
CHEK2	ERBB2*	FANCA	FANCL	FOXA1	MLH1*	MRE11	MSH2*	MSH6*	NBN
PALB2*	PMS2	PTEN*	RAD51B	RAD51C	RAD51D	RAD54L	RB1*	SPOP	TP53*

9 Fusion Genes

ALK	FGFR1	FGFR2	FGFR3	NTRK1	NTRK2	NTRK3	RET	ROS1	
-----	-------	-------	-------	-------	-------	-------	-----	------	--

MSI

*CNV (amplification/deletion) analysis is included for these genes

Biomarkers within the scope of accreditation (ISO15189:2022)

-BRCA1, BRCA2 (SNV/indel detection by NGS)

-MSI (NGS)

Sensitivity: Positive reference standards are tested with the assay, all corresponding mutation sites can be accurately detected, and the positive percent agreement (PPA) for all variants (SNVs, Indels, fusions and CNVs) assessed was 95%.



N° 822

52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 6032138 Fax. (+30) 210 6032148
A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022 (Cert. No 048190009)

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE



Name

-

Report No

26xxxxxxGR

Specificity: Negative reference standards are tested with the assay, and the negative percent agreement (NPA) of SNVs, Indels, fusions and CNVs was 99%.

Limit of Detection (LoD): The detection limit of the method is 1-2% of mutant allelic content, depending on the genomic region. DNA variations detected at frequencies of less than 5% are confirmed using NGS or an alternative method (Real-Time PCR). All fusions detected are confirmed with an alternative method (Real-Time PCR or FISH).

The performance characteristics of the assay are summarized in the following table:

Variant Type	Limit of Detection
Single nucleotide variations (SNV)	VAF $\geq$ 2-5%
Insertions/deletions (Indel)	VAF $\geq$ 2-5%
Fusion (or rearrangement)	VAF $\geq$ 2%
Copy number (CNV)	$\geq$ 3 copies

Disclaimer

1. This test is mainly used to assist clinical decision-making, and the result does not represent clinical decision.
2. The test should be interpreted by combining the actual patient context. The medication information provided only on the basis of genetic test results, and the actual medication should follow the physician's instructions.
3. The clinical trials only present partial relevant clinical recruitment trials. For more comprehensive and updated information, please refer to the website: <https://clinicaltrials.gov/>.
4. As evidence on variants and drugs evolves, previous classifications may later be modified. The interpretation of a variant is based on current available evidence.
5. Sequence variants were reported using Human Genome Variation Society (HGVS) nomenclature. Classification and interpretation of variants follow guidelines of American College of Medical Genetics and Genomics (ACMG), Association of Molecular Pathology (AMP), American Society of Clinical Oncology (ASCO) and College of American Pathologists (CAP).
6. Database and references used: Reference genome (GRCh37), annotation using A Locus Reference Genomic (LRG), database referencing 1000G (phaseIII-ucsc), EXAC (0.3.1), dbSNP (147), PolyPhen2/SIFT (ensdb v73), PhyloP (2013-12-06), Clinvar (2018-8) and Cosmic(V80).

Limitations

1. The test is limited to test genomic variations on DNA level and does not involve RNA level or protein level.
2. Limited tissue detection may not represent the whole DNA variations of lesions because of tumor heterogeneity.
3. Scientific data show that not all patients carry genomic variations that are associated with targeted drug, therefore not all subjects can be matched with targeted therapies or clear resistance mechanism.



52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 6032138 Fax. (+30) 210 6032148
A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022 (Cert. No 048190009)

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE



Name

-

Report No

26xxxxxxGR

4. Genetic variation beyond the detection range of this test or some non-gene mutation related factors such as drug interactions may affect the clinical effects of drugs.
5. The detection method cannot distinguish between somatic mutations and germline mutations effectively without control sample analysis.
6. Every molecular test has an internal 0.5-1% chance of failure. This is due to rare molecular events and factors related to the preparation and analysis of the samples.
7. Improper sample collection, transportation and storage may lead to compromised or invalid results.



52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 6032138 Fax. (+30) 210 6032148
A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022(Cert. No 048190009)

-

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE



Name

-

Report No

26xxxxxxGR

3. Quality Control Results

Quality Control Index	Result	Criterion
Average effective sequencing depth ¹	8400	≥ 500
Tumor cell content ²	30%	≥20%
Overall Assessment ³	PASS	

Note

1. Average effective sequencing depth: Average sequencing depth on target with duplicated reads.
2. An overall tumor cell content percentage of ≥ 20% is recommended for the efficient detection of somatic alterations in the sample analyzed.
3. Overall Assessment: The quality control overall assessment results are divided into two levels: "PASS" and "RISK". When the overall quality assessment result is "RISK", 94-96% of coverage was achieved in the genes analysed, hence there is a small range where clinical actionable variations could be undetected.



52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 6032138 Fax. (+30) 210 6032148
A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022(Cert. No 048190009)

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE



4. Important biomarkers findings

Gene	Finding (VAF)	Gene	Finding (VAF)
<i>ATM</i> mutation	Not Detected	<i>FANCL</i> mutation	Not Detected
<i>ATR</i> mutation	Not Detected	<i>MLH1</i> mutation	Not Detected
<i>BARD1</i> mutation	Not Detected	<i>MRE11</i> mutation	Not Detected
<i>BRAF_V600E</i>	Not Detected	<i>MSI</i> status	MSS
<i>BRCA1</i> mutation	Not Detected	<i>NBN</i> mutation	Not Detected
<i>BRCA2</i> mutation	Not Detected	<i>NTRK1/2/3</i> rearrangement	Not Detected
<i>BRIP1</i> mutation	Not Detected	<i>PABL2</i> mutation	Not Detected
<i>CDK12</i> mutation	Not Detected	<i>RAD51B</i> mutation	Not Detected
<i>CHEK1</i> mutation	Not Detected	<i>RAD51C</i> mutation	Not Detected
<i>CHEK2</i> mutation	Not Detected	<i>RAD51D</i> mutation	Not Detected
<i>ERBB2</i> amplification	Not Detected	<i>RAD54L</i> mutation	Not Detected
<i>FANCA</i> mutation	Not Detected	<i>RET</i> rearrangement	Not Detected

5. Immune Checkpoint inhibitors biomarkers

Biomarker/Variant	Result	Clinical Interpretation
Treatment effect - positive correlation		
<i>MSI</i> status	MSS	-
<i>MLH1</i> suspected germline deleterious mutation	Not Detected	-
<i>MSH2</i> suspected germline deleterious mutation	Not Detected	-
<i>MSH6</i> suspected germline deleterious mutation	Not Detected	-
<i>PMS2</i> suspected germline deleterious mutation	Not Detected	-
<i>KRAS</i> mutation	Not Detected	-
<i>ATM</i> mutation	Not Detected	-
<i>ATR</i> mutation	Not Detected	-
<i>BRCA1</i> mutation	Not Detected	-
<i>BRCA2</i> mutation	Not Detected	-
<i>BRIP1</i> mutation	Not Detected	-
<i>CHEK1</i> mutation	Not Detected	-
<i>CHEK2</i> mutation	Not Detected	-
<i>FANCA</i> mutation	Not Detected	-
<i>MRE11A</i> mutation	Not Detected	-
<i>NBN</i> mutation	Not Detected	-
<i>RAD51B</i> mutation	Not Detected	-



Name		Report No	26xxxxxxGR
RAD51D mutation	Not Detected	-	
RAD54L mutation	Not Detected	-	
TP53 mutation	Detected	May increase the benefit rate of PD-1/PD-L1 inhibitors	
Treatment effect - negative correlation			
ALK rearrangement	Not Detected	-	
PTEN inactivating mutation	Not Detected	-	



52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 60321 38 Fax. (+30) 210 6032148
 A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022(Cert. No 048190009)

-

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE



Name

-

Report No

26xxxxxxGR

6. Interpretations for targeted therapies

Genetic Variation: **TP53: c.817C>T (p.R273C)**

VAF*: 43%

OncoKB

Gene information

TP53 encodes the p53 tumor suppressor protein, a transcription factor that responds to cellular stresses, including DNA damage and oncogenic activation, by inducing downstream anti-tumor responses such as DNA repair and apoptosis (PMID: [11099028](#)). p53 levels are kept low in healthy cells due to negative regulation by MDM2/4, Cop1 and Trim24 and constant degradation by the ubiquitin-proteasome system (PMID: [36859359](#), [36207426](#)). When DNA is damaged, a network of pathways is activated to detect and repair lesions in a cell- and context-specific manner (PMID: [36207426](#)). p53 is rapidly phosphorylated by upstream regulators such as ATM, ATR, and CHL1/2, which results in the accumulation of stable p53 (PMID: [36207426](#)). p53 then binds to specific DNA sequences to direct the expression of a wide variety of genes, including those involved in apoptosis, cell cycle arrest, DNA repair, senescence, stem cell differentiation, autophagy, cellular metabolism, and others (PMID: [27141080](#), [36859359](#), [36207426](#)). Oncogenic mutations of TP53 often result in the dysregulation of p53 function, usually due to structural changes in the DNA binding domain (PMID: [36859359](#)). Loss of p53 function can have various outcomes including tumorigenesis, invasion and metastasis, drug resistance, metabolic reprogramming, immune evasion and overall genomic instability (PMID: [36859359](#)). TP53 is the most commonly mutated gene in human cancers, and germline mutations occur in the cancer predisposition syndrome Li-Fraumeni (PMID: [22713868](#), [21765642](#)). Clinical and preclinical research into drugs that target TP53 is ongoing, notably with MDM2 inhibitors that aim to restore p53 function and are being tested in combination with other cancer therapies (PMID: [36859359](#), [37818252](#)).

Variant Information

The TP53 R273C mutation is likely oncogenic. The TP53 R273C mutation is located in the protein's DNA binding domain. Expression of this mutant in TP53 null mouse embryonic fibroblasts (MEFs) resulted in a significant increase in cell migration relative to TP53 null cells (PMID: [23612969](#)). Structural studies demonstrated that the R273C mutation leads to a dramatic reduction in the DNA binding affinity of p53, although the protein retains wildtype stability (PMID: [23863845](#), [11900253](#)). Expression of this mutation in cell lines that lack TP53 expression did not enhance TP53-mediated transcriptional activity in reporter assays, as compared to expression of wildtype TP53 (PMID: [15037740](#)). Yet, there are no FDA-approved or NCCN-compendium listed treatments specifically for patients with TP53 R273C mutant prostate cancer.

*VAF: Variant Allele Frequency



52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 6032138 Fax. (+30) 210 6032148
A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022 (Cert. No 048190009)

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE



Name

-

Report No

26xxxxxxGR

7. Clinical Trials to consider

TP53 associated clinical trials

[NCT06739395](#)

PHASE2

Title	Precision Medicine Trial Based on Molecular Matching Therapy for Patients With Standard Treatment Exhaustion
Treatment	Olaparib tablet Temozolomide capsule Anlotinib Trametinib tablet Dabrafenib Vebreltinib Enteric Capsules Alpelisib Pill Sacituzumab Govitecan-Hziy 180 MG Lenvatinib Capsules Pazopanib Pill Palbociclib Pill Chidamide PD-1/PD-L1/PD-1&CTLA4 inhibitor Target Gene
Location	China

[NCT06329206](#)

PHASE1

Title	A Phase Ia/Ib Study of GH2616 Tablet in Subjects With Advanced Solid Tumors
Treatment	GH2616 Tablets
Location	China

Press [here](#) for a live search of clinical trials for TP53

52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 60321 38 Fax. (+30) 210 6032148
 A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022(Cert. No 048190009)

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE

Com|P|i|t DX

Name-Report No26xxxxxGR

8. Appendix




Figure 1. Joint consensus recommendation of AMP, ACMG, ASCO and CAP for reporting genetic variants in cancer. [1-2]

1. Li MM, Datto M, Duncavage EJ, et al. J Mol Diagn. 2017 Jan;19(1):4-23.
2. Leichsenring J, Horak P, Kreutzfeldt S, et al. Int J Cancer. 2019 Dec 1;145(11):2996-3010.



Name

-

Report No

26xxxxxxGR

Microsatellite Instability (MSI)

Microsatellite instability (MSI) is a condition of genetic hypermutability resulting from impaired DNA mismatch repair (MMR) mechanisms. It is characterized by the accumulation of mutations in short repetitive DNA sequences known as microsatellites. MSI-H is found across multiple cancer types, most notably in colorectal, endometrial, gastric, and ovarian cancer. It can arise sporadically or as part of Lynch syndrome, a hereditary condition caused by germline mutations in MMR genes (MLH1, MSH2, MSH6, PMS2).

The results of MSI are divided into three types: MSI-H, which means microsatellites are highly unstable; MSS, which means microsatellites are stable; MSI-U, which means that the sample does not meet the MSI evaluation conditions (tissues or pleural fluid samples may not have passed the MSI indicator calculation quality control due to the low DNA and/or content of tumor cells).

Pembrolizumab has received full FDA approval for adult and pediatric patients with unresectable or metastatic MSI-H or dMMR solid tumors (PMID: 35680043, 33264544) — representing the first full approval for an immunotherapy based on a predictive biomarker, regardless of solid tumor type. This approval applies to patients who have progressed after prior treatment and have no satisfactory alternative options. Beyond the tumor-agnostic indication, specific approvals have been granted for tumor types such as endometrial and colorectal cancer.



52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 60321 38 Fax. (+30) 210 6032148
A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022(Cert. No 048190009)

-

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE



Name - Report No 26xxxxxxGR

9. Biomarker findings across all Levels of evidence (A-D)

Biomarker	Result	Therapies with strong clinical significance		Therapies with potential significance		Therapies with potential resistance
		Approved/ Standard of care therapies (Level A)	Therapies in well powered studies (Level B)	Off label/ Investigational therapies (Level C)	Therapies in pre-clinical trials (Level D)	
TP53	Exon 8 c.817C>T (p.R273C) VAF: 43%	-	-	-	-	-



52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 60321 38 Fax. (+30) 210 6032148
A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022(Cert. No 048190009)





Name

-

Report No

26xxxxxxGR

10. References

1. Tops BB, Normanno N, Kurth H, Amato E, Mafficini A, Rieber N, Le Corre D, Rachiglio AM, Reiman A, Sheils O, Noppen C, Lacroix L, Cree IA, Scarpa A, Ligtenberg MJ, Laurent-Puig P. Development of a semi-conductor sequencing-based panel for genotyping of colon and lung cancer by the Onconetwork consortium. BMC Cancer. 2015 Jan 31;15:26. doi: 10.1186/s12885-015-1015-5. (PMID: [25637035](#))
2. <https://civic.genome.wustl.edu/>
3. <http://cancer.sanger.ac.uk/>
4. <https://www.clinicaltrials.gov>
5. <http://atlasgeneticsoncology.org>
6. <https://www.oncokb.org/>
7. <https://www.mycancergenome.org/>
8. <https://pmkb.org/>
9. Cantley LC. **The phosphoinositide 3-kinase pathway.** Science. 2002; 296:1655-7. doi: 10.1126/science.296.5573.1655 (PMID: [12040186](#))
10. Goldstein JL and Wilson JD. **Studies on the pathogenesis of the pseudohermaphroditism in the mouse with testicular feminization.** J Clin Invest. 1972; 51:1647-58. doi: 10.1172/JCI106966 (PMID: [4402348](#))
11. Wang H, et al. **Targeting p53 pathways: mechanisms, structures, and advances in therapy.** Signal Transduct Target Ther. 2023; 8:92. doi: 10.1038/s41392-023-01347-1 (PMID: [36859359](#))
12. Baroni TE, et al. **A global suppressor motif for p53 cancer mutants.** Proc Natl Acad Sci U S A. 2004; 101:4930-5. doi: 10.1073/pnas.0401162101 (PMID: [15037740](#))
13. Addie M, et al. **Discovery of 4-amino-N-[(1S)-1-(4-chlorophenyl)-3-hydroxypropyl]-1-(7H-pyrrolo[2,3-d]pyrimidin-4-yl)piperidine-4-carboxamide (AZD5363), an orally bioavailable, potent inhibitor of Akt kinases.** J Med Chem. 2013; 56:2059-73. doi: 10.1021/jm301762v (PMID: [23394218](#))
14. Georget V, et al. **Trafficking of the androgen receptor in living cells with fused green fluorescent protein-androgen receptor.** Mol Cell Endocrinol. 1997; 129:17-26. doi: 10.1016/s0303-7207(97)04034-3 (PMID: [9175625](#))
15. He B, et al. **Dependence of selective gene activation on the androgen receptor NH2- and COOH-terminal interaction.** J Biol Chem. 2002; 277:25631-9. doi: 10.1074/jbc.M202809200 (PMID: [12000757](#))
16. Ruizeveld de Winter JA, et al. **Androgen receptor expression in human tissues: an immunohistochemical study.** J Histochem Cytochem. 1991; 39:927-36. doi: 10.1177/39.7.1865110 (PMID: [1865110](#))
17. Alaseem AM. **Advancements in MDM2 inhibition: Clinical and pre-clinical investigations of combination therapeutic regimens.** Saudi Pharm J. 2023; 31:101790. doi: 10.1016/j.jsps.2023.101790 (PMID: [37818252](#))
18. Wang Q, et al. **A hierarchical network of transcription factors governs androgen receptor-dependent prostate cancer growth.** Mol Cell. 2007; 27:380-92. doi: 10.1016/j.molcel.2007.05.041 (PMID: [17679089](#))
19. Brown TR, et al. **Deletion of the steroid-binding domain of the human androgen receptor gene in one family with complete androgen insensitivity syndrome: evidence for further genetic heterogeneity in this syndrome.** Proc Natl Acad Sci U S A. 1988; 85:8151-5. doi: 10.1073/pnas.85.21.8151 (PMID: [3186717](#))



Name	Report No
-	26xxxxxxGR
20. Antonarakis ES, et al. Prevalence and Spectrum of AR Ligand-Binding Domain Mutations Detected in Circulating-Tumor DNA Across Disease States in Men With Metastatic Castration-Resistant Prostate Cancer. JCO Precis Oncol. 2024; 8:e2300330. doi: 10.1200/PO.23.00330 (PMID: 38781544) 21. Odell AF, et al. A novel p53 mutant found in iatrogenic urothelial cancers is dysfunctional and can be rescued by a second-site global suppressor mutation. J Biol Chem. 2013; 288:16704-16714. doi: 10.1074/jbc.M112.443168 (PMID: 23612969) 22. Wong CI, et al. Steroid requirement for androgen receptor dimerization and DNA binding. Modulation by intramolecular interactions between the NH2-terminal and steroid-binding domains. J Biol Chem. 1993; 268:19004-12. (PMID: 8360187) 23. Bullock AN and Fersht AR. Rescuing the function of mutant p53. Nat Rev Cancer. 2001; 1:68-76. doi: 10.1038/35094077 (PMID: 11900253) 24. Lubahn DB, et al. Sequence of the intron/exon junctions of the coding region of the human androgen receptor gene and identification of a point mutation in a family with complete androgen insensitivity. Proc Natl Acad Sci U S A. 1989; 86:9534-8. doi: 10.1073/pnas.86.23.9534 (PMID: 2594783) 25. Abuetaf Y, et al. DNA damage response revisited: the p53 family and its regulators provide endless cancer therapy opportunities. Exp Mol Med. 2022; 54:1658-1669. doi: 10.1038/s12276-022-00863-4 (PMID: 36207426) 26. Veldscholte J, et al. A mutation in the ligand binding domain of the androgen receptor of human LNCaP cells affects steroid binding characteristics and response to anti-androgens. Biochem Biophys Res Commun. 1990; 173:534-40. doi: 10.1016/s0006-291x(05)80067-1 (PMID: 2260966) 27. Eldar A, et al. Structural studies of p53 inactivation by DNA-contact mutations and its rescue by suppressor mutations via alternative protein-DNA interactions. Nucleic Acids Res. 2013; 41:8748-59. doi: 10.1093/nar/gkt630 (PMID: 23863845) 28. He WW, et al. A frame-shift mutation in the androgen receptor gene causes complete androgen insensitivity in the testicular-feminized mouse. Nucleic Acids Res. 1991; 19:2373-8. doi: 10.1093/nar/19.9.2373 (PMID: 2041777) 29. Fenton MA, et al. Functional characterization of mutant androgen receptors from androgen-independent prostate cancer. Clin Cancer Res. 1997; 3:1383-8. (PMID: 9815822) 30. Aubrey BJ, et al. Tumor-Suppressor Functions of the TP53 Pathway. Cold Spring Harb Perspect Med. 2016; 6:(unknown pages). doi: 10.1101/cshperspect.a026062 (PMID: 27141080) 31. Steck PA, et al. Identification of a candidate tumour suppressor gene, MMAC1, at chromosome 10q23. 3 that is mutated in multiple advanced cancers. Nat Genet. 1997; 15:356-62. doi: 10.1038/ng0497-356 (PMID: 9090379) 32. Vogelstein B, et al. Surfing the p53 network. Nature. 2000; 408:307-10. doi: 10.1038/35042675 (PMID: 11099028) 33. Suzuki K and Matsubara H. Recent advances in p53 research and cancer treatment. J Biomed Biotechnol. 2011; 2011:978312. doi: 10.1155/2011/978312 (PMID: 21765642) 34. Balbas MD, et al. Overcoming mutation-based resistance to antiandrogens with rational drug design. Elife. 2013; 2:e00499. doi: 10.7554/eLife.00499 (PMID: 23580326) 35. Thompson J, et al. Disrupted amino- and carboxyl-terminal interactions of the androgen receptor are linked to androgen insensitivity. Mol Endocrinol. 2001; 15:923-35. doi: 10.1210/mend.15.6.0647 (PMID: 11376111) 36. Freed-Pastor WA and Prives C. Mutant p53: one name, many proteins. Genes Dev. 2012; 26:1268-86. doi: 10.1101/gad.190678.112 (PMID: 22713868) 37. Georget V, et al. Mechanism of antiandrogen action: key role of hsp90 in conformational change and transcriptional activity of the androgen receptor. Biochemistry. 2002; 41:11824-31. doi: 10.1021/bi0259150 (PMID: 12269826)	



52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 60321 38 Fax. (+30) 210 6032148
A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022(Cert. No 048190009)

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE



Name	Report No
-	26xxxxxxGR
38. Yarbrough WG, et al. A single base mutation in the androgen receptor gene causes androgen insensitivity in the testicular feminized rat. J Biol Chem. 1990; 265:8893-900. (PMID: 2341409) 39. Tan J, et al. Dehydroepiandrosterone activates mutant androgen receptors expressed in the androgen-dependent human prostate cancer xenograft CWR22 and LNCaP cells. Mol Endocrinol. 1997; 11:450-9. doi: 10.1210/mend.11.4.9906 (PMID: 9092797) 40. Schweizer MT and Yu EY. Persistent androgen receptor addiction in castration-resistant prostate cancer. J Hematol Oncol. 2015; 8:128. doi: 10.1186/s13045-015-0225-2 (PMID: 26566796) 41. Shang Y, et al. Formation of the androgen receptor transcription complex. Mol Cell. 2002; 9:601-10. doi: 10.1016/s1097-2765(02)00471-9 (PMID: 11931767) 42. Wilson CM and McPhaul MJ. A and B forms of the androgen receptor are expressed in a variety of human tissues. Mol Cell Endocrinol. 1996; 120:51-7. doi: 10.1016/0303-7207(96)03819-1 (PMID: 8809738)	



52, Spaton Ave., 15344, Gerakas, Greece, G.E.MI. nr: 0007856001000 | info@genekor.com, www.genekor.com | Tel. (+30) 210 6032138 Fax. (+30) 210 6032148
 A company certified with ISO 9001:2015 (Cert. No 041150049) and ISO/IEC 27001:2022(Cert. No 048190009)

George Nasioulas, PhD Molecular Biologist, Scientific Director, Medical ID:26025301255

Document Code: Δ21_Prostate_FFPE